



## Animal Health Products





**ANTIBACTERIAL INJECTABLE PRODUCTS**

Ba-Süfla Solution for Injection	215,8 mg/ml Sülfadimidin sodium, 40 mg/ml Trimethoprim.....	6
Baranisin LA 20% Solution for Injection	200 mg/ml Oxytetracycline .....	6
Bavetaxin Solution for Injection	100 mg/ml Tulathromycin .....	7
Clinexin Suspension for Injection	50 mg/ml Cefiofur .....	7
Ekopen Suspension for Injection	200 mg/ml Procaine Penicillin G, 250 mg/ml Dihydrostreptomycin Sulfate .....	8
Fluflor Solution for Injection	300 mg/ml Florfenicol, 27,4 mg/ml Flunixin meglumine.....	8
Gentadur 10% Solution for Injection	100 mg/ml Gentamicin .....	9
Killoxacin 10% Solution for Injection	100 mg/ml Enrofloxacin .....	9
Killoxacin 5% Solution for Injection	50 mg/ml Enrofloxacin.....	10
Linpectan Solution for Injection	50 mg/ml Lincomycin HCL, 100 mg/ml Spectinomycin Sulfate .....	10
Marboline Solution for Injection	100 mg/ml Marbofloxacin.....	11
Max-til 300 Solution for Injection	300 mg/ml Tilmicosin .....	11
Tay-max 200 Solution for Injection	200 mg/ml Tylosin .....	12
Vetiflor 300 Solution for Injection	300 mg/ml Florfenicol .....	12

**ANTIBACTERIAL INTRAMAMMARY PRODUCTS**

Amoklizon LC Intramammary Suspension	200 mg/5g Amoxicillin, 50 mg/5g clavunilic acid, 10 mg/5g prednisolone .....	13
Multivet LC Intramammary Suspension	100 mg Benzylpenicillin procaine, 100 mg Streptomycin sulfate, 100 mg Neomycin sulfate, 10 mg Prednisolone.....	14

**ANTIBACTERIAL ORAL PRODUCTS**

Ba-Matrix Oral Tablet	1000 mg Sulfadiazine, 200 mg Trimethoprim .....	15
Bapar Powder of Oral Solution	100 mg Paromomycin sulfate .....	15
Baranisin Aqua Oral Powder	755 mg/g Oxytetracycline hydrochloride.....	16
Bavedox 10% Oral Solution	115 mg/ml Doxycycline Hyclate.....	16
Bavedox 50% Powder for Oral Solution	577 mg Doxycycline hyclate (equivalent to 500 mg base) .....	17
Bavedox Oral Tablet	115,388 mg/tablet doxycycline hyclate .....	17
Bavermisin Powder for Oral Solution	509 mg/g Erythromycin .....	17
Killoxacin 20% Oral Solution	200 mg/ml Enrofloxacin .....	18
Killoxacin 80% Powder for Oral Solution	800 mg/kg Enrofloxacin.....	18
Killoxacin Oral Tablet	150 mg/tablet Enrofloxacin .....	18
Megatril Oral Solution	100 mg/ml Enrofloxacin .....	19
SulfaMega Oral Suspension	400 mg/ml Sulfamethoxazole, 80 mg/ml Trimethoprim .....	19
Vetiflor Aqua Oral Powder	500 mg/g Florfenicol .....	20

**ANTIPARASITIC INJECTABLE PRODUCTS**

Anipracit Solution for Injection	56,8 mg/ml Praziquantel .....	21
Babesit Solution for Injection	120 mg of imidocarb dipropionate equivalent to 84.198 mg of imidocarb...22	22
Bavermec Fascio Solution for Injection	10 mg/ml Ivermectin, 100 mg/ml Chlorsulon .....	22
Bavet Klosantel Solution for Injection	50 mg Closantel sodium dihydrate .....	23
Günmek 1% Solution for Injection	10 mg/ml Ivermectin.....	23

**ANTIPARASITIC ORAL PRODUCTS**

Balbend 10% Oral Suspension	100 mg Albendazole .....	24
Balbend-K 300 Oral Tablet	300 mg / 1 tablet Albendazole .....	24
Balbend 1200 Oral Tablet	1200 mg / 1 tablet Albendazole .....	25
Klozanid Oral Tablet	250 mg Oxfendazole, 750 mg Oxclozanide.....	25
Wormtal Oral Tablet	50 mg Praziquantel, 144 mg Pyrantel pamoate, 150 mg Febantel .....	26
Aniprazol KK Oral Tablet	50 mg praziquantel, 500 mg fenbendazole/ 1 tablet.....	26

**ANTIPARASITIC TOPICAL PRODUCTS**

Ba-Cyper 10% Dipp Solution	100 mg/ml Cypermethrin.....	27
Ba-Cyper 2,5% Pour-On Solution	25 mg/ml Cypermethrin .....	27
Ba-tick Pour-On Solution	10 mg Flumethrin .....	28
Baveprin Pour on Solution	5 mg Eprinomectin.....	28
Fibroline SM Cat Spot on Solution	100 mg Fipronil, 120 mg (S)-Methoprene .....	29
Fibroline SM Dog Spot on Solution	100 mg Fipronil, 90 mg (S)-Methoprene.....	29

**CLINIC PRODUCTS**

Atromax Solution for Injection	2 mg Atropine sulphate .....	30
Basilazin 2% Solution for Injection	20 mg/ml Xylazine hydrochloride.....	30
Bavet Menbuton Solution for Injection	100 mg Menbuton .....	31
Furex Solution for Injection	20 mg Furosemid .....	31
Tympatrol Oral Emulsion	% 2.5 Polymerized silicon .....	32

**VITAMIN & MINERAL PRODUCT**

Adened AD3E Solution for Injection	500.000 IU Vitamin A, 75.000 IU Vitamin D3, 50 mg Vitamin E .....	33
Bavse Solution for Injection	60 mg Vitamin E (Tocopherol acetate) /ml, 40 mg Vitamin B1 (Thiamine HCl) /ml, 1 mg Sodium Selenite /ml .....	33
Butafan Solution for Injection	100mg/ml Butaphosphan, 0.05 ml/ml vitamin B12 .....	33
Cal-Mix Solution for Injection	190 mg Ca gluconate, 45 mg Ca glucoheptonate, 60 mg Magnesium Chloride, Butaphosphan 4 mg.....	34
Maxivit-C Solution for Injection	200 mg Ascorbic acid .....	34
Maxivit AD3E Solution for Injection	200.000 IU Vitamin A, 30.000 IU Vitamin D3 , 20 mg/ml .....	35
Nerovitavet Solution for Injection	100 mg/ml Vitamin B1, 10 mg/ml Vitamin B6 .....	36

**ANALGESIC AND ANTI-INFLAMMATORY PRODUCTS**

Ba- Keto Solution for Injection	100 mg/ml Ketoprofen.....	37
Bavet Meloxicam Solution for Injection	5 mg Meloxicam .....	37
Bavet Meloxicam X8 Solution for Injection	40 mg/ml Meloxicam .....	38
Fundamin Solution for Injection	82,95mg Flunixin meglumine equivalent to 50 mg Flunixin .....	38
Metazin Solution for Injection	500 g Metamizole Sodium.....	39









## Ba-Sülfa

Solution for Injection

**COMPOSITION:** 215.8 mg sulfadimidin sodium , 40 mg trimetophrim

**TARGET SPECIES:** cattle, horse, cat, dog

**INDICATIONS:** It is used in the treatment of respiratory system, gastrointestinal system, urogenital system infections, skin and nail infections caused by microorganisms sensitive to sulfadimidine and trimethoprim.

**PACKAGING:** 50 ml, 100 ml, 250 ml

**DOSE:** 1 ml/10-15 kg bodyweight, for cattle use intramuscularly or intravenously, for horse use intravenously only.  
1 ml/15 kg bodyweight for cat and dogs use intramuscularly, intravenously or subcutaneously

**WITHDRAWAL TIME:**

**Meat:** 14 days (cattle)

**Milk:** 5 days



\* Please read the leaflet before use.

## Baranisin LA 20%

Solution for Injection

**COMPOSITION:** Oxytetracycline dihydrate equivalent to 200 mg Oxytetracyclin

**TARGET SPECIES:** cattle, sheep, goat

**INDICATIONS:** It indicated to treatment of systemic, regional and respiratory system infections caused by bacteria sensitive to oxytetracycline.

**PACKAGING:** 50 ml, 100 ml, 250 ml

**DOSE:** 1 ml for 10 kg/bw (i.m)  
If needed the dose is repeated after 4-5 days

**WITHDRAWAL TIME:**

**Meat:** 28 days

**Milk:** 10 days (cattle), 7 days (sheep, goat)



\* Please read the leaflet before use.

# Bavetaxin

## Solution for Injection

**COMPOSITION:** 100 mg/ml tulatromycin

**TARGET SPECIES:** cattle, sheep

**INDICATIONS:** Cattle; It is indicated for the treatment and metaphylaxis of bovine respiratory disease (BRD) caused by Mannheimia haemolytica, Pasteurella multocida, Histophilus somni and Mycoplasma bovis. It is indicated for the treatment of infectious keratoconjunctivitis caused by Morexella bovis sensitive to tulatromycin

**Sheep:** It is indicated for the early treatment of pododermatitis caused by virulent Dichelobacter nodosus

**PACKAGING:** 100 ml, 250 ml

**DOSE:** Cattle: 2,5 mg tulatromycin/kg b.w. (1 ml/40 kg b.w.) single dose via s.c.  
Sheep: 2,5 mg tulatromycin/kg b.w. (1 ml/40 kg b.w.) single dose via i.m. route

**WITHDRAWAL TIME:**

**Meat:**

Cattle: 22 days,

Sheep: 16 days

Not to be used in animals producing milk intended for human consumption or animals expected to give birth in two months that will produce milk for human consumption.



\* Please read the leaflet before use.

# Clinexin

## Suspension for Injection

**COMPOSITION:** ceftiofur hydrochloride equivalent to 50 mg ceftiofur

**TARGET SPECIES:** cattle

**INDICATIONS:** It is used in cattle for the treatment of respiratory and soft tissue infections caused by ceftiofur-sensitive bacteria. It is used in the treatment of Mannheimia sp. (Pasteurella haemolytica), Pasteurella multocida, Respiratory tract infections caused by Haemophilus somnus, Acute interdigital necrobacillosis (fork rot, pododermatitis) caused by Fusobacterium necrophorum and Bacterioides melanogenicus, acute mastitis with postnatal e coli, Arcanobacterium necropoygenium, and Arcanobacterium bacterium acute puerperal metritis.

**PACKAGING:** 50 ml, 100 ml, 250 ml

**DOSE:** 1 ml/50 kg/bw intramuscularly or subcutaneously, one application for every 3-5 days

**WITHDRAWAL TIME:**

**Meat:** 8 days

**Milk:** 0 day



\* Please read the leaflet before use.

# Ekopen

Suspension for Injection

**COMPOSITION:** 200 mg procain penicillin G, 250 mg dihydrostreptomycin sulphate

**TARGET SPECIES:** cattle, sheep, horse, cat and dog.

**INDICATIONS:** Ekopen Injectable Suspension is used in secondary infections caused by respiratory infections, pneumonia, umbilical infections, Listeriosis, Leptospirosis, meningitis, septicemia, urogenital system infections, salmonellosis and viral infections caused by susceptible bacteria in cattle, sheep and horses.

**PACKAGING:** 50 ml, 100 ml, 250 ml

**DOSE:** PVia deep intramuscular injection

**Sheep, cattle, horse:** 1 ml for 25 kg body weight for 3 days

**Cat, dog:** 0,5 ml for 5 kg bodyweight for 3-5 days

**WITHDRAWAL TIME:**

**Meat:** 35 days (cattle, sheep)

**Milk:** 72 hours (6 milkings)



\* Please read the leaflet before use.

# Fluflor

Solution for Injection

**COMPOSITION:** 300 mg florfenicol, 27.4 mg flunixin meglumin equivalent to 16.5 mg flunixin

**TARGET SPECIES:** cattle

**INDICATIONS:** It is used in cattle for the treatment of respiratory infections caused by Mannheimia haemolytica, Pasteurella multocida, Mycoplasma bovis and Histophilus somni.

**PACKAGING:** 100 ml, 250 ml

**DOSE:** 2 ml/15 kg bw/day, one application subcutaneously

**WITHDRAWAL TIME:**

**Meat:** 46 days.

**Milk:** Not authorized for use in lactating animals producing milk for human consumption.



\* Please read the leaflet before use.

# Gentadur 10%

Solution for Injection

**COMPOSITION:** 100 mg gentamicin

**TARGET SPECIES:** cattle, horse, dog, cat

**INDICATIONS:** It is used against gentamicin sensitive bacteria causing respiratory, digestive and urogenital system infections, septicaemia and other soft tissue infections treatment in cattle, horse, dog and cat.

**PACKAGING:** 50 ml, 100 ml, 250 ml

**DOSE:** Cattle: 2 ml / 100kg body weight for 3-7 days  
Horse: 6.5 ml/100kg bw for 3-5 days  
Dog and cat: 0.2ml / 5 kg body weight for 3-5 days

**WITHDRAWAL TIME:**  
Meat: 139 days(cattle), 103 days(calves) for 3 injections,  
192 days for 6 injections, 214 days for 7 injections  
Milk: 7 days



\* Please read the leaflet before use.

# Killoxacin 10%

Solution for Injection

**COMPOSITION:** 100 mg enrofloxacin

**TARGET SPECIES:** cattle, sheep, goat

**INDICATIONS:** It is used for the treatment of sensitive gram negative, gram positive and Mycoplasma related infections in cattle and sheep and goats.

**PACKAGING:** 20 ml, 50 ml, 100 ml, 250 ml

**DOSE:** Practical dose: 1 ml for 20 kg bw/day Product may given subcutaneously or intravenously. Treatment should continue for 3-5 days.

**WITHDRAWAL TIME:**  
Meat: 5 days (cattle i.v.), 12 days (cattle s.c.), 4 days (sheep), 6 days (goats)  
Milk: 3 days (sheep, cattle i.v.), 4 days (goats, cattle s.c.)



\* Please read the leaflet before use.

# Killoxacin 5%

Solution for Injection

**COMPOSITION:** 50 mg enrofloxacin

**TARGET SPECIES:** cattle, sheep, goat, cat, dog

**INDICATIONS:** It is used in the treatment of infections caused by gram-negative and gram-positive bacteria sensitive to enrofloxacin in calves, sheep, goats, cats and dogs.

**PACKAGING:** 20 ml

**DOSE:** practical dosage is 1 ml for 10 kg. Killoxacin 5 % Solution for Injection is administered subcutaneous or intravenously.

**WITHDRAWAL TIME:**

**Meat:** 5 days (cattle i.v.), 12 days (cattle s.c.) 4 day (sheep), 6 days (goats)

**Milk:** 3 days (sheep), 4 days (goats). Do not use in cattle producing milk for human consumption



\* Please read the leaflet before use.

# Linpectan

Solution for Injection

**COMPOSITION:** 50 mg lincomycin, 100 mg spectinomycin sulfate

**TARGET SPECIES:** sheep, lambs, calves, cattle, dog, cat, broiler

**INDICATIONS:** It is used in the treatment of respiratory, digestive, urinary system infections and soft tissue infections and arthritis caused by bacteria sensitive to lincomycin and spectinomycin in target species.

**PACKAGING:** 50 ml, 100 ml, 250 ml

**DOSE:** Sheep-goat: 1 ml for 10 kg bodyweight during 3-5 days Calf

**Steer, Cattle:** 1 ml for 10 kg bodyweight, intramuscularly, daily during 3-5 days (first day 2 injections)

**Dog-Cat:** 1 ml for 5 kg body weight 1-2 a day, during 3-7 days.

**Chickens and Turkey:** 1 ml for 5 kg, subcutaneously for 3 days

**WITHDRAWAL TIME:**

**Meat:** 21 days (cattle), 14 days (sheep and goat)

Should not given to chickens producing eggs for human consumption .

Do not use in lactating sheep and goat producing milk for human consumption



\* Please read the leaflet before use.

# Marboline

Solution for Injection

**COMPOSITION:** 100 mg marbofloxacin

**TARGET SPECIES:** cattle

**INDICATIONS:** It is indicated for respiratory tract infections caused by *Pasteurella multocida*, *Mannheimia haemolytica* ve *Histophilus somni* and used for the treatment of acute mastitis and infections caused by *Escherichia coli* and *Mycoplasma bovis*.

**PACKAGING:** 100 ml

**DOSE:** For respiratory track infections dose is 2 ml/25 kg b.w. (i.m.). For *Mycoplasma bovis* infections and acute mastitis dose is 1 ml/50kg b.w./ day (i.m.,s.c.), should be applied for 3 days.

**WITHDRAWAL TIME:**

**Meat:** 3 days (8 mg/ kg b.w.), 6 days (2 mg/ kg b.w.)

**Milk:** 3 days (6 milkings, 8 mg/ kg b.w.), 2 days (4 milkings, 2 mg/ kg b.w.)



\* Please read the leaflet before use.

# Max-til 300

Solution for Injection

**COMPOSITION:** 300 mg tilmicosin equivalent to tilmycosin phosphate

**TARGET SPECIES:** cattle, sheep

**INDICATIONS:**

**Cattle:** Treatment of bovine respiratory tract infections caused by *Mannheimia haemolytica* and *Pasteurella multocida*, used in the treatment of interdigital necrobacillosis.

**Sheep:** Treatment of bovine respiratory tract infections originating from *Mannheimia haemolytica* and *Pasteurella multocida*, Foot Rot therapy from *Dichelobacter nodosus* and *Fusobacterium necrophorum*, It is used in the treatment of acute mastitis caused by *Staphylococcus aureus* and *Mycoplasma agalactia*.

**PACKAGING:** 50 ml, 100 ml, 250 ml

**DOSE:** Practical dosage: 1 ml for 30 kg Use subcutaneous only.

**WITHDRAWAL TIME:**

**Meat:** 70 days (cattle), 42 days (sheep)

**Milk:** 18 days (sheep) 36 days(cattle)



\* Please read the leaflet before use.



# Tay-max 200

Solution for Injection

**COMPOSITION:** 200 mg tylosin

**PACKAGING:** 50 ml, 100 ml, 250 ml

**TARGET SPECIES:** cattle, sheep, goat, cat, dog

**DOSE:**  
Applied intramuscularly and only applied slow intravenous for cattle.

**INDICATIONS:**Cattle (adult): metritis caused by gram positive bacteria, respiratory system infections, Streptococcus spp., Staphylococcus spp.

Treatment of mastitis caused by Fusobacterium necrophorum and interdigital necrobacillosis caused by Fusobacterium necrophorum, Calf: Necrobacillosis (Fusobacterium necrophorum) and respiratory system infections

**Sheep-goat:** metritis caused by gram positive bacteria, respiratory system infections, Gram positive bacteria or Mycoplasma spp. mastitis caused by

**Dog:** Respiratory tract infections and otitis caused by susceptible bacteria

**Cat:** respiratory tract infections caused by susceptible bacteria

**Cattle:** 2.5-5 ml/100 kg bw

**Sheep-goat:** 2.5 ml/50 kg bw

**Cat-dog:** 0.35-0.55 ml/10 kg bw

**WITHDRAWAL TIME:**

**Meat:** 28 days (cattle) 42 days (sheep, goat)

**Milk:** 108 hours



\* Please read the leaflet before use.

# Vetiflor 300

Solution for Injection

**COMPOSITION:** 300 mg florfenicol

**DOSE:**  
1 ml/ 15 kg b.w. (i.m.), 2 ml/ 15 kg b.w.(s.c. for cattle only) treatment should continue at 48 hours intervals.  
**Sheep:** 1 ml/ 15 kg bw (i.m.) for 3 days

**TARGET SPECIES:** cattle, sheep

**INDICATIONS:**Cattle: Treatment and metaphylaxis of respiratory tract infections caused by florfenicol-susceptible strains of Mannheimia haemolytica, Pasteurella multocida and Histophilus somni. Before metaphylactic use, the presence of disease in the flock should be ensured.

**Sheep:** respiratory tract infections from Mannheimia haemolytica and Pasteurella multocida.

**WITHDRAWAL TIME:**

**Meat:** 30 days (i.m., cattle), 39 days (sheep, 44 days (s.c., cattle) Do not use in lactating cattle producing milk (including dry period) for human consumption

**PACKAGING:** 50 ml, 100 ml, 250 ml



\* Please read the leaflet before use.



# Amoklizon LC

Intramammary Suspension

## COMPOSITION:

Amoxicillin trihidrat equivalent to 200 mg amoxicillin  
Potasyum clavulanate equivalent to 50 mg clavulanic acid  
10 mg Prednisolone

TARGET SPECIES: Cattle

INDICATIONS: Chronic mastitis treatment of lactating cows caused by sensitive microorganisms

## PACKAGING:

5 g tubes 48 or 120 tubes in buckets

DOSE: One syringe is applied to each mammary lobe. 3 applications in every 12 hours.

## WITHDRAWAL TIME:

Meat: 7 days  
Milk: 84 hour (7 milkings)



\* Please read the leaflet before use.



# Multivet LC

Intramammary Suspension

**COMPOSITION:** 100 mg Benzylpenicillin procaine, 100 mg Streptomycin sulfate, 100 mg Neomycin sulfate, 10 mg Prednisolone

**TARGET SPECIES:** cattle

**INDICATIONS:** It is used for the treatment of acute and subacute mastitis caused by bacteria such as Streptococcus spp., Staphylococcus spp., Klebsiella and E.coli which are sensitive to the combination streptomycin, benzylpenicillin and neomycine in lactating cows. It is also used for the prevention of inflammation and pain that occurs with Prednisolone in its content.

**PACKAGING:** 5 g tubes 48 or 120 tubes in buckets

**DOSE:** Infuse one syringe into each affected quarter. Repeat this treatment in 24 hours and apply for 3 consecutive days.

**WITHDRAWAL TIME:**

**Meat:** 7 days (cattle)

**Milk:** 5 days



\* Please read the leaflet before use.



# Ba-Metrix

Oral Tablet

**COMPOSITION:** 1000 mg sulfadiazin 200 mg trimethoprim

**DOSE:** 1 tablet/ 40 kg b.w./ day during 5 days

**TARGET SPECIES:** calves

**WITHDRAWAL TIME:**  
Meat: 15 days

**INDICATIONS:** Ba-Metrix Oral Tablet is mainly indicated for the treatment of bacterial diarrhea in calves, it can also be used in the treatment of acute salmonellosis and bacterial pneumonia.

**PACKAGING:** 2x 5 tablets



\* Please read the leaflet before use.

## Baranisin Aqua

**COMPOSITION:** 755 mg Oxytetracyclin HCl

**TARGET SPECIES:** Salmon, Rainbow trout

**INDICATIONS:** BARANİSİN AQUA Oral Powder is indicated for the treatment of the following diseases of the fresh water and salt water fishes:

- Frunculosis caused by *Aeromonas salmonicida* subsp.
- Yersiniosis caused by *Yersinia ruckeri*
- Columnaris caused by *Flexibacter columnaris*

**PACKAGING:** 1 kg, 5 kg

**DOSE:** Practical dose is 75 mg/ 10 kg b.w.  
It is used for 4 days.

**WITHDRAWAL TIME:** Until daily heat total reaches 720°C

Oral Powder



\* Please read the leaflet before use.

## Bavedox 10%

**COMPOSITION:** 115 mg/ml doxycycline hyclate

**TARGET SPECIES:** Calves, Chicken, Turkey

**INDICATIONS:** It is indicated to treatment of respiratory and digestive system infections on calves, broilers and turkeys

**PACKAGING:** 100 ml, 1 L

**DOSE:** Recommended dose of product is mixed in water  
**Calves;** 0.08 ml product/b.w./day via oral route during 3-5 days  
**Chicken;** 0.21 ml product/b.w./day via oral route during 3-5 days  
**Turkey;** 0.25 ml product/b.w./day via oral route during 3-5 days

**WITHDRAWAL TIME:**

**Meat:** 7 days (calves), 5 days (chicken), 12 days (turkey)

**Egg:** Do not use in animals producing egg for human consumption.

**Milk:** Do not use in animals producing milk for human consumption.

Not to be used within 4 weeks before egg production period.

Oral Solution



\* Please read the leaflet before use.

## Bavedox 50%

Powder of Oral Solution

**COMPOSITION:** 577 mg Doxycycline hyclate (equivalent to 500 mg base)

**TARGET SPECIES:** Non-ruminating Calves, Chicken, Turkey

**INDICATIONS:** It is indicated to treatment of respiratory and digestive system infections on calves, broilers and turkeys

**PACKAGING:** 1 kg

**DOSE:**

Calves; 17.3 mg product/b.w./day) via oral route during 3 or 5 days

Chicken; 43.3 mg product/b.w./day via oral route during 3 or 5 days

Turkey; 50.2 mg product/b.w./day) via oral route during 3 or 5 days

**WITHDRAWAL TIME:**

**Meat:** 7 days (calves), 5 days (chicken), 12 days (turkey)

**Egg:** Do not use in animals producing egg for human consumption.

**Milk:** Do not use in animals producing milk for human consumption.

Not to be used within 4 weeks before egg production period.



\* Please read the leaflet before use.

## Bavedox

Oral Tablet



**COMPOSITION:** 115,388 mg 7 tablet doxycycline hyclate  
**TARGET SPECIES:** Dogs

**INDICATIONS:** It is indicated for treatment of respiratory system infections such as rhinitis, tonsillitis caused by the doxycycline-sensitive microorganisms

\* Please read the leaflet before use.

**PACKAGING:** 10 tablets (1x10) or 30 tablets (3x10) blister packaging in cardboard box

**DOSE:**  
1 tablet/10 kg bw for 5 days

**WITHDRAWAL TIME:**  
Not applicable

## Bavermisin

Powder for Oral Solution

**COMPOSITION:** 550 mg erythromycin thiocyanate equivalent to 509 mg erythromycine

**TARGET SPECIES:** Chicken, Turkey

**INDICATIONS:** Treatment of infections associated with Mycoplasma

**PACKAGING:** 1 kg

**DOSE:** It is used as 20-40 mg/g erythromycine base/day via oral route during 3 or 5 days  
Practically 39-79 g/1000 kg b.w.

**WITHDRAWAL TIME:**

**Meat:** 3 days (chicken), 6 days (turkey)

**Egg:** 5 days



\* Please read the leaflet before use.



# Killoxacin 20%

Oral Solution

**COMPOSITION:** 200 mg enrofloxacin

**TARGET SPECIES:** Chicken, Turkey, Rabbit

**INDICATIONS:** It is used in the treatment of respiratory and gastro intestinal system infections caused by susceptible bacteria in chicken, turkey and rabbits.

**PACKAGING:** 1L, 2.5L

**DOSE:** Practically 5 ml of KILLOXACIN 20 % Oral Solution per 100 kg body weight is added to drinking water daily. It is used for 3-5 days (5 days in rabbit)

**WITHDRAWAL TIME:**  
**Meat:** 7 days (chicken), 13 days (turkey), 3 days (rabbit)

**Egg:** Do not use in animals producing egg for human consumption. Not to be used within 14 days to egg laying period.



\* Please read the leaflet before use.

# Killoxacin

Oral Tablet

**COMPOSITION:**  
 150 mg/tablet enrofloxacin

**TARGET SPECIES:**  
 Dogs

**INDICATIONS:**  
 It is indicated to treat following infections. In dogs; single or combined respiratory system, lower urinary tract (single or with prostatitis), upper urinary tract infections caused by Escherichia Coli or Proteus mirabilis, skin infections, superficial and deep pyoderma, otitis externa caused by gram positive and gram negative bacteria .

**PACKAGINGS:**  
 10 tablets (1x10) or 30 tablets (3x10) blister packaging in cardboard box

**DOSE:**  
 1 tablet/30 kg bw for 10-21 days depending on the severity and type of infection.

**WITHDRAWAL TIME:**  
 Not applicable



\* Please read the leaflet before use.

# Killoxacin 80%

Powder for Oral Solution

**COMPOSITION:** 800 mg/g enrofloxacin

**TARGET SPECIES:** chicken, turkey, rabbit

**INDICATIONS:** It is used for the treatment of infections caused by the above listed sensitive bacteria in chicken, turkey and rabbits.

**Chickens:** Mycoplasma gallisepticum, Mycoplasma synoviae, Avibacterium paragallinarum, Pasteurella multocida

**Turkey:** Mycoplasma gallisepticum, Mycoplasma synoviae, Pasteurella multocida

**Rabbit:** E. coli and Pasteurella multocida

**PACKAGING:** 1 kg

**DOSE:**

**Chickens and turkeys:** 10 mg enrofloxacin/kg bodyweight per day for 3-5 consecutive days. Treatment period is 5 consecutive days in mixed infections and chronic progressive forms.

**Rabbits:** 10 mg/kg bodyweight per day for 5 consecutive days.

**WITHDRAWAL TIME:**

**Meat:** 7 days (chicken), 13 days (turkey), 3 days (rabbit)

**Egg:** Do not use in animals producing egg for human consumption. Not to be used within 14 days to egg laying period.



\* Please read the leaflet before use.

# Megatril

**COMPOSITION:** 100 mg enrofloxacin

**TARGET SPECIES:** chicken, turkey, rabbit

**INDICATIONS:** It is indicated for the treatment of the infections caused by gram negative and positive bacteria and Mycoplasma spp. susceptible to enrofloxacin in broiler turkeys, chickens and rabbits.

**PACKAGING:** 100 ml, 3L, 5L

**DOSE:** The daily dose to be administered is 10 mg/kg body weight by adding drinking water for 3-5 days (5 days for rabbit).

**WITHDRAWAL TIME:**

**Meat:** 7 days (chicken), 13 days (turkey), 3 days (rabbit) Should not be given to chickens and turkeys producing eggs for human consumption, not to be used within 14 days to egg laying period.

Oral Solution



\* Please read the leaflet before use.

# SulfaMega

**COMPOSITION:** 400 mg sulfamethoxazol, 80 mg trimethoprim

**TARGET SPECIES:** poultry

**INDICATIONS:** It is indicated for the treatment and prevention of Escherichia coli respiratory tract infections susceptible to trimethoprim and sulfamethoxazole in diseased herds.

**PACKAGING:** 100 ml, 1L, 2.5L

**DOSE:** There should be 1 ml Sulfamega Oral Suspension per 3.64 kg body weight. Treatment should be continued for 3-4 days

**WITHDRAWAL TIME:**

**Meat:** 10 days Should not given to poultry producing eggs for human consumption

Oral Suspension



\* Please read the leaflet before use.



# Vetiflor Aqua

Oral Powder

**COMPOSITION:** 500 mg florfenicol

**TARGET SPECIES:** aquaculture

**INDICATIONS:** VETIFLOR AQUA Oral Powder is indicated for the treatment of the diseases of the fresh water (trout and carp) and salt water fishes (sea bream, sea bass, Salmon) caused by gram negative and positive bacteria.

**PACKAGING:** 1 kg, 5 kg

**DOSE:** Practical dosage: 0.02 g/kg bodyweight by adding fish feed for 10 days

**WITHDRAWAL TIME:**

**Meat:** 15 days (5°C) , 7 days (10°C)



\* Please read the leaflet before use.



# Anipracit

Solution for Injection

**COMPOSITION:** 56,80 mg paraziqantel

**TARGET SPECIES:** Cats, dogs

**INDICATIONS:** Treatment of cestod infestations in cats and dogs

**PACKAGING:** 20 ml

**DOSE:**

Practically, 0.1 ml is injected for 1 kg body weight. (1 ml/10 kg b.w.) (s.c.)(i.m.)

**WITHDRAWAL TIME:** Not applicable



\* Please read the leaflet before use.



# Babesit

Solution for Injection

**COMPOSITION:** 120 mg imidocarb dipropionate

**TARGET SPECIES:** cattle, horse, dog

**INDICATIONS:** It is indicated for the treatment and prevention of babesiosis in cattle, dog and horse.

**PACKAGING:** 20 ml, 50 ml

**DOSE:**

**Cattle:** 1 ml/100 kg bw for treatment, 2.5 ml/100 kg bw for prevention (i.m.)(s.c.)

**Dog:** 0.25 ml/10 kg bw for treatment, 0.5 ml/10 kg bw for prevention (i.m.)(s.c.)

**Horse:** 2 ml/100 kg bw for treatment and prevention (i.m.)(s.c.)

**WITHDRAWAL TIME:Meat:** 213 days

**Milk:** 21 days



\* Please read the leaflet before use.

# Bavermec Fascio

Solution for Injection

**COMPOSITION:** 10 mg ivermectin, 100 mg clorsulon

**TARGET SPECIES:** cattle

**INDICATIONS:** It is used in the prevention and treatment of gastrointestinal nematodes, lung nematodes (adult and 4th stage larvae), subcutaneous nematode, eye nematode, nocturnal agents (larval stage), lice, scabies and liver trematodes in cattle.

**PACKAGING:** 50 ml, 100 ml, 250 ml

**DOSE:** 1 ml for 50 kg/ bodyweight

Use subcutaneously only

**WITHDRAWAL TIME:**

**Meat:** 66 days,

Not authorised to use on cattle that produce milk for human consumption.

Not to be used within 60 days before giving birth.



\* Please read the leaflet before use.

# Bavet Klosantel

Solution for Injection

**COMPOSITION:** Closantel sodium dihydrate equivalent to 50 mg Closantel sodium base

**TARGET SPECIES:** cattle, sheep

**INDICATIONS:** Treatment of closantel sensitive trematods, nematods and arthropod larvae invasions.

**PACKAGING:** 100 ml

**DOSE:**

1-2 ml/20 kg b.w., s.c. For cattle, max. 20 ml (6 ml in calves) for an injection area.  
1-2 ml/20 kg b.w., s.c. for in sheep, max. 5 ml for an injection area.

**WITHDRAWAL TIME:**

**Meat:** 84 days (cattle), 107 days (sheep).

Not to be used on animals producing milk for human consumption or cattle at last trimester.



\* Please read the leaflet before use.

# Günmek %1

Solution for Injection

**COMPOSITION:** 10 mg ivermectin

**TARGET SPECIES:** cattle, sheep

**INDICATIONS:** It is used in the treatment and control of gastro-intestinal nematodes, lung nematodes, subcutaneous nematodes, eye nematode, noca agents, scabies, lice and tick infestations in cattle.

It is used in the treatment and control of gastro-intestinal nematodes, lung nematodes, nasal nematodes, scabies and lice infestations in sheep.

**PACKAGING:** 50 ml, 100 ml, 250 ml

**DOSE:** Practical dosage is 1 ml for 50 kg bodyweight

Only given subcutaneously.

**WITHDRAWAL TIME:**

**Meat:** 49 days (cattle), 42 days (sheep) Not to be used within 60 days to birth in animals producing milk intended for human consumption



\* Please read the leaflet before use.

# Balbend 10%

Oral Suspension

**COMPOSITION:** 100 mg albendazole

**TARGET SPECIES:** cattle, sheep, goat

**INDICATIONS:** It is indicated to treatment and prevention of infestations caused of gastrointestinal nematode, lungworm and trematodes.

**PACKAGING:** 1 L

**DOSE:**

**Cattle:** 7,5 ml/ 100 kg b.w. for gastrointestinal roundworms and lungworms and 10 ml/ 100 kg b.w. for adult flukes

**Sheep:** 1 ml/ 20 kg b.w. for gastrointestinal roundworms and lungworms and 1,5 ml/ 20 kg b.w. for adult flukes, Dicrocoeliasis dose is 3 ml/20 kg b.w.

**Goats, worm and fluke** dose is 1,5 ml/ 20 kg b.w. and Dicrocoeliasis dose is 6 ml/40 kg b.w.

**WITHDRAWAL TIME:**

**Meat:** 14 days for cattle, 10 days for sheep and goats

**Milk:** 3 days for cattle, 6 days for sheep and goats (not for human consumption during treatment)



\* Please read the leaflet before use.

# Balbend-K 300

Oral Tablet

**COMPOSITION:** 300 mg albendazole

**TARGET SPECIES:** sheep, cattle, goat

**INDICATIONS:** It is indicated to treatment and prevention of infestations caused by worms, liver flukes.

**PACKAGING:** 10 tablets

**DOSE:**

**Cattle:** 1 tablet/40 kg b.w. for gastrointestinal roundworms and lungworms, and 1 tablet/30 kg b.w. for adult flukes

**Sheep:** 1/2 tablet/30 kg b.w. for gastrointestinal roundworms and lungworms, and 1 tablet/40 kg b.w. for adult flukes, Dicrocoeliasis dose is 2 tablet/40 kg b.w.

**Goat:** 1 tablet/40 kg b.w. for gastrointestinal roundworms and lungworms and 2 tablet/40 kg b.w. for Dicrocoeliasis

**WITHDRAWAL TIME:**

**Meat:** 14 days for cattle, 10 days for sheep and goat

**Milk:** 3 days for cattle, 6 days for sheep and goat (not for human consumption during treatment)



\* Please read the leaflet before use.

# Balbend 1200

Oral Tablet

**COMPOSITION:** 1200 mg albendazole

**TARGET SPECIES:** cattle

**INDICATIONS:** It is indicated to treatment and prevention of infestations caused by worms, liver flukes.


**PACKAGING:** 10 tablets

**DOSE:** 1 tablet/ 160 kg b.w. for worms and yung flukes and 1 tablet/ 120 kg b.w.for adult flukes

**WITHDRAWAL TIME:**

**Meat:** 14 days

**Milk:** 3 days (6 milkings) (not for human consumption during treatment)



\* Please read the leaflet before use.

# Klozanid

Oral Tablet

**COMPOSITION:** oxfendazol 250 mg, oxiclozanid 750 mg

**TARGET SPECIES:** cattle, sheep

**INDICATIONS:** KLOZANİD Oral Tablet is used in prophylaxis and treatment of gastrointestinal and pulmonary infestation by mature and larvae of pinworm, and mature liver flukes, mature Moniezia sp. in cattle and sheep

- a) Liver Flukes
- b) Gastrointestinal pinworms  
Stomach  
Intestine  
Large intestine
- c) Pulmonary pinworms
- d) Tapeworm

**PACKAGING:** 10, 20, 50 tablet

**DOSE:**

**Lamb:** ½ tablet for 25 kg

**Sheep:** 1 tablet for 50 kg

**Calf-Steer:** 2 tablets for 100 kg

**Cattle:** 10 tablet for 500 kg

**WITHDRAWAL TIME:**

**Meat:** 28 days (cattle) , 21 days (sheep)

**Milk:** 5 days (cattle) Do not use in lactating sheep producing milk for human consumption



\* Please read the leaflet before use.



# Wormtal

Oral Tablet

**COMPOSITION:** 50 mg praziquantel, 144 mg pirantel pamoat, 150 mg febantel

**TARGET SPECIES:** dog

**INDICATIONS:**The following names of dogs are used to control and control parasitic invasions caused by nematode and cestode species. *Toxocara canis*, *Toxascaris leonina*(adults and species close to becoming adults)

*Uncinaria stenocephala*, *Ancylostoma caninum* (adults)

*Trichuris vulpis*

*Echinococcus granulosus*, *Echinococcus multilocularis*, *Dipylidium caninum*, *Taenia* sp.

**PACKAGING:** 50 tablet, 1x50 sachets

**DOSE:**Practical dosage for dogs: 1 tablet for 10 kg bodyweight

Tablets may also be administered half according to the weight of the animals.

**WITHDRAWAL TIME:**Not applicable



\* Please read the leaflet before use.

# Aniprazol KK

Oral Tablet

**COMPOSITION:** 50 mg praziquantel, 500 mg febendazole

**DOSE:**1 tablet for 10 kg bodyweightTablets may also be administered half according to the weight of the animals

**TARGET SPECIES:** Cat and dog

**WITHDRAWAL TIME:** Not applicable

**INDICATIONS:** Used for the treatment of infestations caused by roundworms and tapeworms in dogs and cats

**PACKAGING:** 3\*10 tablets blister1\*50 tablets sachet in cardboard box



\* Please read the leaflet before use.

# Ba-Cyper 10%

Dipp Solution

**COMPOSITION:** 100 mg/ml cypermethrin

**DOSE:** 1 lt product per 500 lt bath solution

**TARGET SPECIES:** Sheep

**WITHDRAWAL TIME:**

**Meat:** 14 days

Not for use in sheep producing milk for human consumption.

**INDICATIONS:** It is indicated to prevention and treatment of blowfly strike, lice, keds and ticks on sheep It is indicated to prevention and treatment of sheep scab caused by pyrethroid sensitive mites.

**PACKAGING:** 100 ml, 500 ml, 1000 ml



\* Please read the leaflet before use.

# Ba-Cyper 2,5%

Pour-On Solution

**COMPOSITION:** 25 mg/ml cypermethrin

**DOSE:**

Pharmacologic dosage: 5-10 mg/kg b.w.

**TARGET SPECIES:** Cattle

**WITHDRAWAL TIME:**

**Meat:** 10 days

**Milk:** "0" days

**INDICATIONS:** It is indicated to control of lice and flies in cattle

**PACKAGING:** 100 ml, 500 ml



\* Please read the leaflet before use.



## Ba-tick

Pour-On Solution

**COMPOSITION:** 10 mg flumethrin

**TARGET SPECIES:** cattle, sheep

**INDICATIONS:** Ba-tick pour-on solution is used for the control and treatment against biting flies, chewing and biting lice (*Haematopinus eurysternus*, *Linognathus vituli*, *Bovicola bovis*, *B. ovis*, *Melophagus ovinus*), ticks, mange mites (scabies) and other ecto parasites related infections in cattle and sheep

**PACKAGING:** 100 ml, 500 ml

**DOSE:** The application dose is 1 mg/kg for tick control in cattle and sheep, and 2 mg/kg body weight for scabies and lice control.

**WITHDRAWAL TIME:**

**Meat:** 21 days (sheep) 5 days (cattle)

**Milk:** 8 days

Not authorised to use on sheep that produce milk for human consumption.



\* Please read the leaflet before use.

## Baveprin

Pour-On Solution

**COMPOSITION:** 5 mg eprinomectin

**TARGET SPECIES:** cattle

**INDICATIONS:** It is used for the treatment and control of gastrointestinal pinworms (adult and larval forms), lung pinworms (adult and larval forms), lice, chorioptic and sarcoptic mange and *Haematobia irritans* in cattle.

**PACKAGING:** 500 ml

**DOSE:** 1 ml/10 kg body weight. It should be applied along the back line from the shoulders to the head of the tail. Its effect lasts 2-4 weeks

**WITHDRAWAL TIME:** Meat: 10 days, Milk: "0" days



\* Please read the leaflet before use.

# Fibroline SM Cat

Spot on Solution

**COMPOSITION:** 100 mg Fipronil, 120 mg (S)-Methoprene

**PACKAGING:** 0.5 tubes

**TARGET SPECIES:** cat

**DOSE:** A tube or pipette is placed on the skin and gently squeezed several times to completely drain the contents into a single drop on the skin.

**INDICATIONS:**It is used in the treatment and prevention of the following external parasite infestations in cats:

- Elimination of fleas
- Elimination of ticks
- Elimination of biting lice

**WITHDRAWAL TIME:**Not applicable



\* Please read the leaflet before use.

# Fibroline SM Dog

Spot on Solution

**COMPOSITION:** 100 mg Fipronil, 90 mg (S)-Methoprene

**PACKAGING:** 0,67 ml-1,34 ml-2,68 ml-4,02 ml tubes

**TARGET SPECIES:** dog

**DOSE:** A tube or pipette is placed on the skin and gently squeezed several times to completely drain the contents into a single drop on the skin.

**INDICATIONS:**It is used in the treatment and prevention of the following external parasite infestations in dogs:

- Elimination of fleas
- Elimination of ticks
- Elimination of biting lice

Can be used as a support for the control of Fleas Allergic Dermatitis (FAD)

0.67 ml tube/2-10 kg bw  
1.34 ml tube/10-20 kg bw  
2.68 ml tube/20-30 kg bw  
4.02 ml tube/30-40 kg bw

**WITHDRAWAL TIME:** Not applicable



\* Please read the leaflet before use.

# Atromax

Solution for Injection

**COMPOSITION:** 2 mg atropine sulphate

**PACKAGING:** 20 ml

**TARGET SPECIES:** Horse, cat, dog

**DOSE:** Subcutaneously (s.c.) intramuscular(i.m.), intravenous (i.v.)  
0.25-5 mg/kg body weight

**INDICATIONS:** For stimulation of hearth movements in horse, dog and cats, at cases of pain, bladder, uretra and kidney spasms colics and constrictions as spasmolytic, at organic phosphorus poisoning with aracoline, pilocarpin and neostigmin poisonings as antidote at general anesthesia for reduction of saliva, for prevention of vago-vagal reflexes and vagal tonus negative effects on hearth as preanesthetic medication.

**WITHDRAWAL TIME:** Not applicable



\* Please read the leaflet before use.

# Basilazin 2%

Solution for Injection

**COMPOSITION:** Xylazine hydrochloride 20 mg/ml

**PACKAGING:** 50 ml

**TARGET SPECIES:** cattle, horse, cat, dog

**DOSE:** Dosing varies depending on the degree of sedation required and the type of animal. See the prospectus for detailed information.

**INDICATIONS:** Basilazin 2% is a sedative with analgesic and muscle relaxant properties for use in cattle, horses, dogs and cats in cases where sedation is required.

**WITHDRAWAL TIME:**

**Meat:** 1 day (cattle)

**Milk:** 0 (zero) day



\* Please read the leaflet before use.

# Bavet Menbuton

Solution for Injection

**COMPOSITION:** 100 mg menbuton

**TARGET SPECIES:** cattle, sheep, dog, horse

**INDICATIONS:** Stimulation of hepato-digestive activity in case of digestive disorders and hepatic insufficiency.

**PACKAGING:** 100 ml

**DOSE:**

**Cattle:** 1 ml for 15-20 kg/ bodyweight by intravenously

**Sheep, goat and calves:** 1 ml for 10 mg/kg bodyweight by deep intramuscular injection or intravenous slowly

**Horse:** 1 ml for 20-40 kg/ bodyweight by intravenous injection

**WITHDRAWAL TIME:** '0' days for meat and milk



\* Please read the leaflet before use.

# Furex

Solution for Injection

**COMPOSITION:** 20 mg Furosemid

**TARGET SPECIES:** cattle, horse, cat, dog

**INDICATIONS:** Furex Solution for Injection is used in order for elimination of fluids, accumulating and causing the formation of edema due to the disorders in circulatory system, liver, kidneys and any other reasons, from the tissues, body cavities, joints, tendons and surfaces through the diuresis, and in order for acceleration of diuresis in case of intoxications, and also it is used in case of emergencies such as acute lung and brain edema.

**PACKAGING:** 20 ml, 50 ml

**DOSE:** Practical dosage: 2,5 - 5 ml/100 kg/bw for cattle and horses (slowly intravenous), 1,25 -2,5 ml/10 kg/bw for dog and cats ( slowly intravenous or intramusculary )

**WITHDRAWAL TIME:** "0 days" meat and milk (i.v.)



\* Please read the leaflet before use.



# Tympatrol

**COMPOSITION:** 2.5% polymerized silicon

**TARGET SPECIES:** cattle, horse, sheep, goat

**INDICATIONS:** It is indicated for acute and chronic meteorismus (foamy, gaseous bloating) in cattle, sheep, goats and horses, and for rumen bloating observed in septicemia in lambs, calves and goat kids.

**PACKAGING:** 100 ml

**DOSE:** TYMPATROL, used orally or intraruminally.

**Cattle - Horse:** 50-75 ml

**Calves - Foal:** 15-25 ml

**Steer:** 25-50 ml

**Sheep - Goat:** 15-25 ml

**WITHDRAWAL TIME:** "0" days for milk and meat

Oral Emulsion



\* Please read the leaflet before use.

# Adened AD3E

Solution for Injection

**COMPOSITION:** 500.000 IU Vitamin A, 75.000 IU Vitamin D3, 50 mg Vitamin E

**TARGET SPECIES:** cattle, sheep, goat, rabbit

**INDICATIONS:** A, D3 and E avitaminosis and treatment for related diseases and malnutritions.

**PACKAGING:** 100 ml, 250 ml

**DOSE:**

**Cattle:** 5-10 ml (i.m.)

**Calves:** 2-4 ml (i.m.)

**Sheep, Goat:** 1-2 ml (i.m.)

**Lamb, Kid, Rabbit:** 0,5 ml (s.c.)

**WITHDRAWAL TIME:**

**Meat:** 28 days

**Milk:** 5 days



\* Please read the leaflet before use.



500.000 IU / ml



75.000 IU / ml



50 mg / ml

# Butafan



Solution for Injection

**COMPOSITION:** 100 mg Butaphosphan, 0,05 mg vitamin B12 (cyanocobalaminum)

**TARGET SPECIES:** cattle, horse, cat, dog

**INDICATIONS:** It is used as a supplement in muscle, metabolism or reproductive system disorders where the use of phosphorus and cyanobalamin is required in cattle, horse, cats and dogs.

**PACKAGING:** 50 ml, 100 ml, 250 ml

**DOSE:**

**2-5 ml/ 100 kg b.w. (i.v.)** in horses and cattle.

**0,35-0,6 ml/ 10 kg b.w.** in foals and calves.

**0,25-2,5 ml/ 10 kg b.w. (i.v., i.m., s.c.)** in dogs.

**0,5-5 ml/ 5 kg b.w. (i.v., i.m., s.c.)** in cats. Doses must be split in half with 1-2 weeks delay for chronic conditions in cats and dogs.

**WITHDRAWAL TIME:** '0' days for meat and milk (cattle, horse)



\* Please read the leaflet before use.

# Bavse

Solution for Injection

BAVSE Solution for Injection is used for white muscle disease, myopathies, growth delay due to selenium and Vit E deficiency, and nutritional-related loss of fertility in lambs and calves, cows, sheep and foals.

**COMPOSITION:** 60 mg Vitamin E (Tocopherole acetate), 40 mg Vitamin B1 (Thiamin HCl), 1 mg Sodium Selenite in each ml.

**PROPHYLAXIS:** Depending on the dose, repeat after 1 week if necessary.

**TREATMENT:** A maximum of 4 doses are administered, at 2 weeks interval.

**PACKAGING:** 100 ml

**WITHDRAWAL TIME:** Meat: 28 days.

Not to be used in animals producing milk intended for human consumption

Species	Body weight (kg)	Dose	
		Prophylaxis	Therapeutic
Cattle, horse ( After the 5th month of pregnancy )	100	5-10 ml	-
Sheep, goat ( After the 3rd month of pregnancy )	50	1-2 ml	-
Calf, Foal	50	0,5-1 ml	1,2 ml
Lamb,goat (1-7 days)	10	0,25 ml	0,25-0,5 ml
Lamb,goat (7-30 days)	10	0,25-0,5 ml	0,5-1,0 ml
Lamb, goat (older than 1 month)	20	0,5-1,0 ml	1,2 ml



# Cal-Mix

Solution for Injection

**COMPOSITION:**

190 mg Ca gluconate  
45 mg Ca glucoheptonate  
60 mg Magnesium Chloride  
4 mg Butaphosphan

**TARGET SPECIES:** cattle, sheep, horse, goat, dog

**INDICATIONS:** It is indicated for the protection from and treatment of acute and chronic metabolism disorders resulted from the deficiency of calcium, magnesium and phosphor in cattle, horses, sheep, goats and dogs.

**PACKAGING:** 100 ml, 250 ml

**DOSE:** Use intramuscularly, intravenously or subcutaneously  
Practical dosage: 1 ml/1 kg bw

**WITHDRAWAL TIME:** "0" days for meat and milk (cattle, sheep, goat)



\* Please read the leaflet before use.

# Maxivit-C

Solution for Injection

**COMPOSITION:** 200 mg ascorbic acid

**TARGET SPECIES:** cattle, sheep, horse, goat, cat, dog

**INDICATIONS:** It is used to treatment and prevention diseases caused by ascorbic acid (vitamin C) deficiency in animals. It is used in clinical and supportive treatment in cases of developmental and adaptation deficiencies, anemia, hypocalcemia, mycotoxiosis and intoxication

**PACKAGING:** 50 ml, 100 ml

**DOSE:**

**Cattle:** 7 - 12 ml /day i.m , s.c , i.v twice a week during 3-6 weeks  
**Steer:** 2 - 4 ml/day i.m, s.c, i.v twice a week during 3 weeks  
**Sheep:** 2 - 4 ml/day i.m , s.c twice a week during 3 weeks  
**Goat:** 2 - 4 ml/day i.m , s.c twice a week during 3 weeks  
**Horse:** 10 -20 ml /day i.m, i.v  
**Cat:** 0,3 - 0,6 ml/day i.m, i.v  
**Dog:** 0,3 – 2,5 ml/day i.m, i.v

**WITHDRAWAL TIME:** "0" days for meat and milk



\* Please read the leaflet before use.





## Maxi-Vit AD<sub>3</sub>E

### COMPOSITION:

Vitamin A Propionat	200,000 I.U. (80 mg)
Vitamin D3	30.000 I.U (0,7875 mg)
Tocopherol Acetate	20 mg

**TARGET SPECIES:** cattle, sheep, goat, rabbit

**INDICATIONS:** MAXIVIT AD<sub>3</sub>E Injectable Solution is indicated for to prevent deficiencies and treatment of diseases due to avitaminosis of these vitamins in cattle, sheep, goat and rabbits.

**PACKAGING:** 100 ml

### DOSE:

Cattle:	5-10 ml
Calves:	2-4 ml
Sheep- Goat:	1-2 ml
Lamb-rabbit:	0.5 ml

### WITHDRAWAL TIME:

Meat:	28 days
Milk:	5 days

Solution for Injection



\* Please read the leaflet before use.



# Nerovitavet

Solution for Injection

**COMPOSITION:** 100 mg vitamin B1 (tiamin hcl) ,  
10 mg vitamin B6 (piridoxin hcl)

**TARGET SPECIES:**cattle, sheep, goat, cat, dog

**INDICATIONS:**NEROVITAVET Solution for Injection is used against Vitamin B1 (Thiamin HCl) and Vitamin B6 (Pyridoxine HCl) deficiencies or as supportive to treatment in cattle, sheep, goat, cat and dog.

**PACKAGING:**100 ml

**DOSE:**

**Cattle - Horse:** 400 kg 10 -20 ml daily  
**Steer - Heifer:** 200 kg 5 – 10 ml daily  
**Sheep - Goats:** 100 kg 2,5 - 5 ml daily  
**Cat - Dog:** 5 kg 1 -2 ml daily

**WITHDRAWAL TIME:**“0” days for milk and meat



\* Please read the leaflet before use.

# Ba-Keto

Solution for Injection

**COMPOSITION:** 100 mg ketoprofen

**TARGET SPECIES:** cattle, horse

**INDICATIONS:**

**Horses:**

Diseases affecting the osteoarticular and muscular-skeletal system associated with acute pain and inflammation

- Lameness of traumatic origin
- Arthritis
- Osteitis, spavin
- Tendinitis, bursitis
- Naviculitis
- Laminitis
- Myositis

**Cattle:**

- alleviation of pain (e.g. from pressure trauma) resulting from parturient paresis;

- reduction of the pyrexia and distress associated with bacterial respiratory disease when used in conjunction with antimicrobial therapy as appropriate;
- improvement of the recovery rate in acute clinical mastitis, including acute endotoxin mastitis, caused by gram negative micro-organisms, in conjunction with antimicrobial therapy;
- alleviation of pain associated with udder oedema following calving.
- reduction of pain associated with musculoskeletal disorders
- after calving

**PACKAGING:** 100 ml

**DOSE:**

**Cattle:** 3 ml/ 100 kg b.w. use intramuscular or intravenously, for up to 3 days

**Horse:** 1 ml/ 45 kg b.w. for horse use intravenously, for 3-5 days.

**WITHDRAWAL TIME:**

**Meat:** 4 days

**Milk:** Zero days.



\* Please read the leaflet before use.

# Bavet Meloxicam

Solution for Injection

**COMPOSITION:** 5 mg meloxicam

**TARGET SPECIES:** cattle, dog, cat

**INDICATIONS:** In cattle; It is used together with appropriate antibiotics to reduce the clinical manifestations of acute respiratory tract infection. In dogs; It is used to relieve inflammation and relieve pain in acute and chronic musculoskeletal diseases, and to reduce inflammation and pain after orthopedic and soft tissue operations. in cats; It is used to reduce post-operative pain after ovariohysterectomy and soft tissue operations.

**PACKAGING:** 20 ml, 50 ml, 100 ml

**DOSE:**

**Cattle:** 10 ml/100 kg bw subcutaneously, intramuscularly or intravenously or intravenously

**Dogs:** 0.4 ml/10 kg bw subcutaneously

**Cat:** 0.12ml/2 kg bw subcutaneously

**WITHDRAWAL TIME:**

**Meat:** 15 days (Cattle)

**Milk:** 5 days (10 milkings) (cattle)



\* Please read the leaflet before use.

# Bavet Meloxicam X8

Solution for Injection

**COMPOSITION:** 40 mg meloxicam

**TARGET SPECIES:** cattle, horse

**INDICATIONS:** It is used together with appropriate antibiotics to reduce the clinical manifestations of acute respiratory infection.

**PACKAGING:** 20 ml, 100 ml

**DOSE:**

**Cattle:** 1.25 ml / 100 kg b.w. by single dose subcutaneously, intramuscularly or intravenously (sc) or intravenous (iv) injection.

**Horse:** It is administered by a single dose intravenous (iv) i 1.5 ml / 100 kg b.w.

**WITHDRAWAL TIME:**

**Meat:** 15 days (Cattle)

**Milk:** 5 days (10 milkings) (cattle)



\* Please read the leaflet before use.

# Fundamin

Solution for Injection

**COMPOSITION:** 82,95mg flunixin meglumine equivalent to 50 mg flunixin

**TARGET SPECIES:** cattle, horse

**INDICATIONS:**  
**Horse:** Painful and inflammatory conditions of the skeletal-muscular system and colic.  
**Cattle:** Accompanying proper treatment, to reduce clinical symptoms formed during pulmonary infections.

**PACKAGING:** 50 ml, 100 ml, 250 ml

**DOSE:** Intravenous application should be slow and administered at body temperature.

2 ml/ 50 kg b.w. (slow i.v., i.m.) for 1-3 days in cattle.

1 ml/ 50 kg b.w. (slow i.v.) for 1-5 days in horses.

**WITHDRAWAL TIME:**

**Meat:** 31 days (i.m.), 10 days (i.v.)

**Milk:** 36 hours (i.m.), 24 hours (i.v.)



\* Please read the leaflet before use.



# Metazin

Solution for Injection

**COMPOSITION:** 500 g metamizole sodium

**TARGET SPECIES:**cattle, horse, dog

**INDICATIONS:**Metazine injectable solution is used in the symptomatic treatment of the following cases in cattle, horses and dogs.

As analgesic, Spasmolithically, Antipyretically, Antirheumatic

**PACKAGING:** 100 ml

**DOSE:** It is administered intramuscularly or slow intravenously.

**Cattle:** 16-32 ml/400 kg bw

**Steer-Heifer:** 4-8 ml/100 kg bw

**Horse:** 20-50 ml/50 kg bw

**Foal:** 2-5 ml/50 kg bw

**WITHDRAWAL TIME:**

**Meat:** 12 days Do not use in lactating cows producing milk for human consumption



\* Please read the leaflet before use.



İstanbul Tuzla Organize Sanayi Bölgesi (İTOSB)  
12. Cadde No:8 34959  
Tepeören-Tuzla / İstanbul - TÜRKİYE

T: +90 0216 593 37 40  
F: +90 0216 593 37 41  
info@bavet.com.tr  
www.bavet.com.tr